



# Combivac C<sup>®</sup>

## Freeze dried live attenuated Newcastle Disease Virus + Infectious Bronchitis Virus Vaccine

Newcastle disease is caused by a paramyxovirus. Signs of disease in birds include hemorrhage and inflammation of respiratory and intestinal tracts and occasional nervous signs. Infectious bronchitis is caused by a coronavirus.

Signs of disease in chickens include inflammation of respiratory and urogenital systems. Combination vaccine for both infections is for efficient easy application.

### COMPOSITION:

Freeze dried live attenuated vaccine against Newcastle disease and infectious bronchitis in chickens. Each dose contains at least  $10^{6.0}$  EID<sub>50</sub> Newcastle Disease Virus strain Clone and at least  $10^{3.0}$  EID<sub>50</sub> Infectious Bronchitis Virus strain H120.

### INDICATIONS:

Combivac C is indicated for vaccination of poultry against Newcastle disease virus and infectious bronchitis virus infections.

### DOSAGE AND ADMINISTRATION:

One dose per bird.

#### Administration:

Intraocular / intranasal route:

- Dissolve the vaccine in physiological saline solution (35 ml).
- Administer one drop of the vaccine using a dropper into one eye or nostril.
- Make sure that the drop of vaccine is inhaled by the bird.
- Use the vaccine immediately after reconstitution (within two hours in cool conditions).

Oral route via drinking water:

- Reconstitute the vaccine with

cold pure drinking water, free of antiseptics, and add to it powdered skimmed milk (2 g/L) to stabilize the vaccine's activity.

- The drinking water temperature containing the vaccine must not exceed 15-20 °C.
- It is advisable to deprive the birds of water 1-2 hours prior to vaccination.
- The vaccine should be given in the early morning or the cool period on a hot day.
- The vaccine should be diluted in adequate quantity of final drinking water in order be consumed within one to two hours after dissolving the vaccine in water according to age of the birds and climatic conditions as follows:  
1-7 days of age  
Dilute 1000 doses in 5 liters  
1-2 weeks of age  
Dilute 1000 doses in 10 liters  
4-8 weeks of age  
Dilute 1000 doses in 20 liters  
Above 8 weeks  
Dilute 1000 doses in 40 liters

Aerogenic route via aerosol / spraying.

- Spray the vaccine solution over the birds, at a distance of 30 to 40 cm, using a coarse sprayer producing.
- Close windows, fans and heaters while spraying.
- The recommended amount of distilled water per 1000 doses is as follows:  
1-7 days of age      200 ml  
2 weeks of age      300 ml  
4 weeks of age      400 ml

### WITHDRAWAL PERIOD:

Zero days.

### CONTRAINDICATIONS:

Do not vaccinate sick or stressed birds.

### SPECIAL PRECAUTIONS:

- Deprive birds of drinking water 1-2 hours before vaccination.
- Apply usual aseptic procedures.
- Use only clean, antiseptic or disinfectant-free materials for the preparation of the vaccine solution.
- The vaccine should be administered to the birds within 2 hours period after reconstitution.
- Unused opened vaccine containers and waste material should be decontaminated with suitable disinfectants (phenolics, iodophores or formaldehydes) at the recommended concentration.

### STORAGE AND SHELF LIFE:

- Store between +2 °C to +8 °C.
- Protect from sunlight.
- The vaccine remains stable for 2 years from production date stated on the label under the recommended storage conditions.

### PRESENTATION:

Vials of: 1000 doses and 2500 doses.

### FOR VETERINARY USE ONLY.



Jovac

#### JORDAN BIOINDUSTRIES CENTER

Tel. +962 6 5232162, Fax: +962 6 5232210

P.O. Box 43, Amman 11941- Jordan

E-mail: info@jovaccenter.com, www.jovaccenter.com